

# Continuous behavioural 'switching' in human spermatozoa and its regulation by Ca<sup>2+</sup>-mobilising stimuli

Achikanu, Cosmas; Correia, Joao; Guidobaldi, HA; Giojalas, LC; Barratt, Christopher L. R.; Da Silva, S M; Publicover, Steve

DOI:  
[10.1093/molehr/gaz034](https://doi.org/10.1093/molehr/gaz034)

License:  
Creative Commons: Attribution (CC BY)

Document Version  
Peer reviewed version

Citation for published version (Harvard):  
Achikanu, C, Correia, J, Guidobaldi, HA, Giojalas, LC, Barratt, CLR, Da Silva, SM & Publicover, S 2019, 'Continuous behavioural 'switching' in human spermatozoa and its regulation by Ca<sup>2+</sup>-mobilising stimuli', *Molecular Human Reproduction*, vol. 25, no. 8, gaz034, pp. 423-432. <https://doi.org/10.1093/molehr/gaz034>

[Link to publication on Research at Birmingham portal](#)

## Publisher Rights Statement:

Checked for eligibility: 21/06/2019  
Cosmas Achikanu, Joao Correia, Hector Guidobaldi, Laura Giojalas, Christopher L R Barratt, Sarah Martins Da Silva, Stephen Publicover, Continuous behavioural 'switching' in human spermatozoa and its regulation by Ca<sup>2+</sup>-mobilising stimuli, MHR: Basic science of reproductive medicine, , gaz034, <https://doi.org/10.1093/molehr/gaz034>

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## **Continuous behavioural ‘switching’ in human spermatozoa and its regulation by Ca<sup>2+</sup>-mobilising stimuli**

Cosmas Achikanu<sup>1,2</sup>, Joao Correia<sup>1</sup>, Hector Guidobaldi<sup>3,4</sup> Laura Giojalas<sup>3,4</sup> Christopher L.R. Barratt<sup>5,6</sup>, Sarah Martins Da Silva<sup>5,6</sup> & Stephen Publicover<sup>1,7\*</sup>,

<sup>1</sup>School of Biosciences, University of Birmingham, Birmingham, UK.

<sup>2</sup>Current address: Department of Applied Biochemistry, Faculty of Applied Natural Sciences, Enugu State University of Science and Technology, Ebeano City, PMB 01660, Enugu, Enugu State, Nigeria.

<sup>3</sup>Centro de Biología Celular y Molecular, Facultad de Ciencias Exactas, Físicas y Naturales (FCEFN), Universidad Nacional de Córdoba (UNC), Córdoba, Argentina

<sup>4</sup>Instituto de Investigaciones Biológicas y Tecnológicas, Consejo de Investigaciones Científicas y Técnicas (CONICET), UNC-FCEFN, Córdoba, Argentina

<sup>5</sup>Reproductive and Developmental Biology, School of Medicine, Ninewells Hospital and Medical School,

<sup>6</sup>University of Dundee, Dundee DD19SY, UK 4Assisted Conception Unit, Ninewells Hospital Dundee, Dundee DD19SY, UK

<sup>7</sup>Centre for Human Reproductive Science, University of Birmingham, UK

\*Correspondence address: s.j.publicover.ac.uk



## **Abstract**

Human sperm show a variety of different behaviours (types of motility) that have different functional roles. Previous reports suggest that sperm may reversibly switch between these behaviours. We have recorded and analysed the behaviour of individual human sperm (180 cells in total), each cell monitored continuously for 3-3.5 min either under control conditions or in the presence of  $\text{Ca}^{2+}$ -mobilising stimuli. Switching between different behaviours was assessed visually (1 s bins using 4 behaviour categories), and was verified by fractal dimension analysis of sperm head tracks. In the absence of stimuli, approximately 90% of cells showed at least one behavioural transition (mean rate under control conditions =  $6.4 \pm 0.8$  transitions.min<sup>-1</sup>). Type 1 behaviour (progressive, activated-like motility) was most common but the majority of cells (>70%) displayed at least three behaviour types. Treatment of sperm with  $\text{Ca}^{2+}$ -mobilising agonists had negligible effects on the rate of switching but increased the time spent in type 2 and type 3 (hyperactivation-like) behaviours ( $P < 2 \times 10^{-8}$ ; chi square). Treatment with 4-aminopyridine under alkaline conditions ( $\text{pH}_o = 8.5$ ), a highly-potent  $\text{Ca}^{2+}$ -mobilising stimulus, was the most effective in increasing the proportion of type 3 behaviour, biasing switching away from type 1 ( $P < 0.005$ ) and dramatically extending the duration of type 3 events ( $P < 10^{-16}$ ). Other stimuli, including 300 nM progesterone and 1% human follicular fluid, had qualitatively similar effects but were less potent. We conclude that human sperm observed in vitro constitutively display a range of behaviours and regulation of motility by  $[\text{Ca}^{2+}]_i$ , at the level of the single cell, is achieved not by causing cells to adopt a 'new' behaviour but by changing the relative contributions of those behaviours.

**Key words:** spermatozoa – behaviour – motility – calcium - pH

## **Introduction**

The ability of mammalian sperm to change behaviour is essential, as different types of motility are suited to overcoming the different barriers that the sperm encounters as it ascends the female tract. For instance, sperm from mice that are null for CatSper (the primary  $\text{Ca}^{2+}$  channel of sperm) fail to hyperactivate and consequently cannot fully ascend the female tract (as they cannot escape after binding the oviduct wall) and are unable to penetrate the zona pellucida (Carlson, et al., 2003, Ho, et al., 2009). During semen analysis, the motility of a sperm population is typically assessed by estimating quantitative characteristics of the population (% motile, % progressively motile). Use of computer-assisted sperm analysis (CASA) provides more detailed information by tracking the movement of each sperm head, typically for a fraction of a second. With these data, the kinematics of each sperm (usually several hundred cells) are calculated, providing an overview of the nature and variation of motility in the population (Mortimer, 2000). In both cases, the quantitative data are calculated/estimated from a 'snapshot' sample, effectively assuming that each cell has a motility type and that the distribution of these types reflects the nature of the population (Pacey, et al., 1997).

However, when mammalian (including human) sperm are monitored for a longer periods (1-10 s), it becomes apparent that the behaviour of individual cells may change rapidly. For instance, hyperactivated-like motility may occur as intermittent 'bursts' interspersed with activated (progressive) swimming (Cooper, et al., 1979; Katz and Yanagimachi, 1980; Johnson, 1981; Burkman, 1984; Mortimer and Swan, 1995; Pacey, et al., 1995). This alternation of behaviour may well be functionally significant, possibly enhancing penetration of the zona pellucida (Bedford, 1998; Katz et al., 1989) or facilitating detachment from the oviduct wall.

Thus, although it is likely that a primary ‘aim’ of stimuli that affect sperm motility is simply to increase the fraction of the population in which a required behaviour occurs, more subtle effects might be achieved at the level of individual cells. For instance, by biasing the probability of switching towards (or away from) a particular behaviour, or by selectively changing the duration of a specific behaviour, the relative amounts of time spent in each behaviour may be regulated. It is also possible that the absolute duration of an individual period of a specific behaviour is functionally important, such as during interaction with the female tract or oocyte vestments.

We hypothesised that behavioural switching and expression of multiple behaviour types is a common feature of human sperm, even under unstimulated conditions. To better understand both the occurrence of multiple patterns of behaviour in individual human sperm and the way in which behavioural switching is regulated, we have: (i) recorded and analysed the activity of individual cells over a prolonged period (>3 minutes) and (ii) investigated whether this behaviour is modified through elevation of  $[Ca^{2+}]_i$ .

## **Materials and methods**

### **Materials**

All chemicals were obtained from Sigma-Aldrich (Poole, UK) unless stated otherwise. Progesterone (P4) and 4-aminopyridine (4-AP) were prepared as stocks in dimethylsulphoxide (DMSO) at 10 mM and 500 mM respectively. Human follicular fluid (FF) was thawed and diluted in saline on the day of use. Thimerosal was prepared as 100 mM stock in deionised water. Working concentrations were made by diluting in supplemented Earle's balanced salt solution (sEBSS) at the appropriate pH prior to use. DMSO in P4 and 4-AP experiments was 0.003% and 0.4% respectively. 1% DMSO had no detectable effect on kinematic properties of sperm motility (Achikanu et al., 2018).

### **Salines**

sEBSS contained (mM) 90 NaCl, 1.017 NaH<sub>2</sub>PO<sub>4</sub>, 5.4 KCl, 0.81 MgSO<sub>4</sub>, 5.5 C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>, 2.5 C<sub>3</sub>H<sub>3</sub>NaO<sub>3</sub>, 19 CH<sub>3</sub>CH(OH)COONa, 25 NaHCO<sub>3</sub>, 1.8 CaCl<sub>2</sub>, 25 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES; pH 7.4), 0.3% BSA. Osmolality was then adjusted to 291-294 mOsm as necessary by adding NaCl. For buffering at pH 8.5, HEPES was replaced with [tris(hydroxymethyl)methylamino]propanesulfonic acid (TAPS).

### **Selection and preparation of spermatozoa**

Semen samples were from donors with normal sperm concentration and motility (WHO 2010). Samples were obtained by masturbation after 2-3 days sexual abstinence. After liquefaction (30 min), sperm were swum up into sEBSS (60 min), adjusted to  $\approx 6$  million/ml and incubated under capacitating conditions (sEBSS at pH 7.4; 37°C, 6% CO<sub>2</sub>) for 5 hours (Alasmari et al., 2013a).

## **Human follicular fluid**

Follicular fluid (FF) was prepared as described in Brown et al. (2017). Briefly, oocytes (metaphase II) were retrieved by transvaginal aspiration. FF without blood contamination was centrifuged (2500 g for 10 min) to separate cellular components. The supernatant (0.22 µm filtered) was stored (at -20°C) until use.

## **Ethical approval**

Written consent was obtained in accordance with the Human Fertilisation and Embryology Authority (HFEA) Code of Practice (version 8) under local ethical approval (University of Birmingham ERC 07-009 and ERN-12-0570) and (13/ES/0091) from the East of Scotland Research Ethics Service REC1.

## **Long-duration sperm tracking**

Observation chambers, 20 µm-deep, were constructed by mounting a 22x32 mm coverslip (0.13-0.16 mm thickness; Academy, Beckenham, UK) onto a 22x50 mm coverslip (thickness no. 1.5; VWR International). Soda lime glass beads (18-22 µm; Cospheric, Santa Barbara, USA) were dispersed in vacuum grease (Dow-Corning, USA) and small dabs were placed at each corner of the 22 mm x 32 mm coverslip, providing a ≈20 mm space when the two coverslips were pressed together.

Capacitated cells were pelleted by centrifugation and re-suspended in 100 µl aliquots (cell density adjusted to 1.2 million/ml) in sEBSS pH 7.4 or 8.5 as required. The stimulus (1% DMSO control, 300 nM progesterone, 1% FF (containing ≈300 nM P4 (Brown et al. 2017)), 1 µM thimerosal or 2 mM 4-AP) was added to the cell suspension and incubation (37°C) was continued for a further 1-2 min. A 5 µl aliquot of cell suspension was carefully loaded into the coverslip chamber, which was then placed on the platform of a thermostatic controller (TC-



324B, Warner, USA) set at 36.5°C on the stage of an inverted microscope (Nikon Eclipse TE200) equipped with a motorised stage (Prior Scientific, UK). Images were acquired with a sCMOS camera (Zyla 5.5, ANDOR TM Technology, Belfast, UK) through a 10x phase contrast objective.

Imaging was started approximately 5 min after stimulus application. Images were acquired at 50 Hz, 15 ms exposure for a period of 180-216 s using Micro Manager software v1.4.22 (image 1392 x 1040 pixels, 16 bit dynamic range, global camera shuttering). Illumination intensity was reduced as far as possible and applied via a red filter to prevent direct effects of illumination on sperm motility (Shahar, et al., 2011; Gabel, et al., 2018). For each recording, a randomly selected sperm was followed. When the cell approached the edge of the field of view, it was moved back using the motorised stage. Using playback of videos, sperm behaviour was analysed in 1 s blocks, allocating each block to one of four different behaviours (motility types 1, 2, 3 and 4; Table 1; Fig. S1; see results). Observation of videos gave no indication that stage movements induced or modified behavioural switching (Fig. S2). We analysed the relationship between the number of stage movements during a 3 min recording (x) and number of behavioural switches in that 3 min (y). There was no significant relationship ( $y=0.08x+18$ ;  $P=0.9$ ;  $R^2=0.001$ ). Furthermore, in many cells in which switching was frequent, no stage movements were required. We also assessed whether the behaviour changes over the period of the recording might occur if evaporation of saline affected osmolality or pH of the droplet. Neither the rate of switching nor the behaviour score showed a significant trend over the duration of the recording ( $P=0.83$ ;  $P=0.58$  respectively; Fig. S3).

In a subset of experiments, where the head centroid could be reliably detected throughout the recording, the multidimensional motion analysis application in MetaMorph® (Molecular Devices, USA) was used to generate positional information for the sperm head for the entire

recording period. Separate sections of the sperm track (between stage re-centring) were ‘stitched’ together by compensating the offset of ‘x’ and ‘y’ co-ordinates in Microsoft Excel.

## CASA

Hyperactivation was assessed as described previously (Achikanu et al.; 2018). After addition of the stimulus, cells were loaded into pre-warmed 20 µm CASA chamber (36°C). Motility was assessed (approximately 300 s after stimulus application) with a Hamilton Thorn CEROS CASA system (version 14.0). Criteria for hyperactivation were VCL (curvilinear velocity)  $\geq 150 \mu\text{m/s}$ , LIN (linearity)  $< 50\%$  and ALH (amplitude of lateral head movement)  $\geq 7 \mu\text{m}$  (Mortimer, 2000).

## Data analysis

Fractal dimension (FD) was determined from x, y coordinates over a period of 1 s (50 points) for each track point along the trajectory. FD was calculated using the equation

$$FD = \log(n) / [\log(n) + \log(l_d/cd)]$$

where n is the number of points in the interval analysed (50 points),  $l_d$  is the linear distance between the first and the last point and cd is the length of the trajectory (Mortimer et al. 1996). FD was calculated with a homemade macro (paNoel 1.0; Universidad Nacional de Córdoba; <http://www.iibyt.conicet.unc.edu.ar/software/>) using Fiji software (Schindelin et al., 2012). Maximum FD was set at 2.0.

Behaviour score (average of all the recorded scores for each 1 s period over the entire recording), % time (% of total recording period spent in each behaviour type), switching rate (transitions.min<sup>-1</sup>), behaviour dwell time (period during which a single behaviour type occurred) and transition type (behaviour type entered upon a transition) were calculated in Microsoft Excel and statistical analysis was carried out using Minitab18 and Excel.

185 Behaviour score, % time and switching rate were calculated separately for each cell and a  
186 mean was calculated from these values (n=number of cells analysed). Dwell times and  
187 transition types were analysed using all individual behaviour events of the specified type  
188 (n=total number of events). All summary values given in the text are mean  $\pm$  SEM unless  
189 stated otherwise. Before analysis, data were tested for normality (Anderson-Darling) and data  
190 sets were then compared using Chi-square, Student's t-test, Mann-Whitney, 2-way ANOVA  
191 and Kruskal-Wallis (with post-hoc Holm-Bonferroni sequential correction; Gaetano, 2013) as  
192 appropriate. Autocorrelation assessment of periodicity was carried out in Minitab 18.

193

## Results

Using five conditions (control and four different stimuli), each at two different values of pH<sub>o</sub> (7.4 and 8.5), we collected video recordings from a total of 180 cells. By visual analysis of video playback, we categorised behaviour into four types. Types 1, 2 and 3 resembled the patterns of motility normally referred to as activated, transitional and hyperactivated (Figure S1, Table 1; Videos1 and 2; Mortimer, 2000). An additional type of behaviour was occasionally observed, in which the mid-piece and anterior flagellum became tightly curved (forming a 'J' shape or coil) and arrested or 'twitched' (Video 3, Table 1). Since behaviour of this type is seen in cells treated with Ca<sup>2+</sup>-ionophore, where it apparently reflects excess Ca<sup>2+</sup>-loading which takes the cells 'beyond' hyperactivated motility (Sanchez-Cardenas et al., 2018), this was designated type 4 behaviour. This behaviour was typically interspersed with very brief (100-500 ms) bursts of flagellar activity such that the cell became virtually immobilised for up to 20 s (Figure S1d, Video 3).

### Free-swimming sperm cells repeatedly 'switch' behaviour

Examination of the behaviour of cells incubated under control conditions (sEBSS; pH 7.4) showed that the majority (16/18 cells) changed their behaviour at least once during the recording. Type 1 behaviour was seen in 18/18 cells, type 3 occurred in 16/18 (no significant difference in incidence;  $P=0.15$ ; Chi square) and 13 of these cells showed three or (in one case, Figure 1a) all four types of behaviour. Among the 18 control cells analysed, the number of behavioural transitions varied widely (from 0 to 45), with the mean rate being  $6.4 \pm 0.8$  transitions.min<sup>-1</sup>. Dwell-times (length of period during which a single behaviour type occurred) for types 1, 2 and 3 events all formed negative exponential distributions (Figure S4a), with the median duration of type 1 events being significantly greater than those for types 2, 3 and 4 ( $P < 5 \times 10^{-5}$ ; Kruskal-Wallis; Figure 2a). Analysis of transition types showed

that switching between behaviours was not random ( $P < 10^{-10}$ ; Chi square), with transitions into type 4 behaviour being very rare (4.3%) and into type 1 behaviour occurring most frequently (44%; Figure 2c column 1). Consequently, cells spent most time ( $65 \pm 6\%$ ) in type 1 behaviour, more than double that of any of the other behaviour types (Figure 2b column 1;  $P < 0.0005$ ; Kruskal-Wallis with post-hoc comparison). Occurrence of type 4 behaviour was very rare, making up less than 3% of behaviour events and of total time (Figure 2b). Neither the rate of switching nor the type of behaviour expressed showed a significant trend over the recording period ( $P = 0.83$ ;  $P = 0.58$  respectively; Figure S3).

In most cells where behavioural switching occurred there was no clear pattern, as type 1 behaviour was interrupted by 'bursts' of type 2 and type 3 activity (Figure 1a). In a small number of cells (4/18), behaviour appeared to oscillate but analysis by autocorrelation showed only one instance where an oscillatory frequency emerged (Figure S5, compared to Figure 1) and even in this case, the autocorrelation failed to reach significance ( $P > 0.05$ ; Figure S5b). For 3 of the 18 control cells, where video quality was adequate for reliable detection of the sperm head throughout the 3 min recording, we were able to reconstruct entire tracks lasting over 3 min, which clearly showed the observed transitions in behaviour (Figure 1c). To confirm that visual analysis identified genuine behavioural transitions, we used the track coordinates to assess temporal variation of fractal dimension (FD; a numerical value between 1 and 2, which reflects the complexity of the sperm track; Mortimer, et al., 1996). Colour coding of track points for FD confirmed that high values occur at periods of complex motility and overlays of FD and visual analyses confirmed that the same behavioural transitions were identified (Figure 1b,c, Figure S5c,d, Figure S6). Autocorrelation of the FD data in Figure S5 produced near identical results to those obtained through visual analysis (Figure S5b,e). When data from all tracked cells were analysed, the mean FD and

mean behavioural score (average score over the entire recording) for each cell were clearly correlated ( $R^2=0.56$ ;  $P=5*10^{-6}$ ;  $n=28$ ; Figure S7).

### **Manipulation of $[Ca^{2+}]_i$ modifies the pattern of behavioural switching**

A pivotal mechanism for regulation of behaviour in human sperm is  $[Ca^{2+}]_i$  (Publicover, et al., 2007; Suarez, 2008). To investigate the effect of  $[Ca^{2+}]_i$  elevation, we monitored the activity of cells exposed to four different stimuli (300 nM P4, 1% FF [containing approximately 300 nM P4], 2 mM 4-AP and 1 mM thimerosal), all of which are known to increase  $[Ca^{2+}]_i$  and stimulate hyperactivation in human and other sperm, but with varying potency (Ho and Suarez, 2001; Alasmari, et al., 2013b; Brown, et al., 2017; Achikanu et al., 2018). CASA assessments, carried out during the period of behavioural data collection, using cells prepared under identical conditions, confirmed the efficacy and relative potency of these stimuli (Figure S8a). The proportion of cells in which we observed switching between behaviours was not changed by any of these manipulations (Table 2), but the relative incidence (% time) of the different behaviour types was significantly modified ( $P<2*10^{-8}$ ; chi square). Incidence of type 1 behaviour was reduced by all four stimuli ( $P<0.05$ ) whereas 4-AP significantly increased the proportion of type 3 behaviour ( $P=2*10^{-6}$ ) and thimerosal significantly increased the occurrence of both types 2 and 3 behaviours ( $P<0.05$ ; Kruskal-Wallis with post-hoc comparison; Figure 2b). Examination of switching showed that this change was partly due to biasing of switching towards types 2 and 3 behaviours in stimulated cells (P4  $P=0.1$ ; FF  $P<0.05$ ; 4-AP and thimerosal  $P<5*10^{-5}$ ; Chi-square; Figure 2c), but there were also marked effects on dwell times of the different behaviours. Type 1 events were significantly briefer in stimulated cells (except for those exposed to thimerosal; Figure 2d) whereas type 2 and type 3 events were markedly prolonged in cells exposed to 4-AP and to thimerosal (Figure 2e,f; Figure S6), such that the overall frequency of behaviour switching

was significantly lower than in controls ( $P=0.02$  and  $P=0.0004$  for 4-AP and thimerosal respectively; Kruskal-Wallis).

### **Effect on behaviour of elevated $pH_o$**

To investigate the effect on behavioural switching of elevated  $pH_o$ , sperm were prepared using the standard protocol but then re-suspended in saline buffered at pH 8.5, which causes stable alkalinisation of the cytoplasm (increase from 6.9 to 7.2) within 200-300 s and increases levels of hyperactivated motility (Achikanu et al., 2018; Figure S8a). The proportion of sperm in which switching occurred was not significantly changed at elevated pH except in cells exposed to 4-AP, where switching was observed in only 10/20 cells ( $P=0.005$  compared to 4-AP at pH 7.4; Table 2). The frequency of switching at  $pH_o=8.5$ , compared to cells incubated at  $pH_o=7.4$ , was consistently lower ( $P<2*10^{-7}$ , 2-way ANOVA; Figure 3a), but when the five conditions were analysed separately, this effect of pH was significant only in cell exposed to FF ( $P<0.01$ ) and 4-AP ( $P<0.001$ ; Mann-Whitney). Similarly to our observations at  $pH_o=7.4$ , at  $pH_o=8.5$  all of the stimuli significantly changed the relative abundance (% time) of the four behaviours (Figure 3b;  $P<10^{-14}$ ; Chi-square), with the most striking effect being in 4-AP-treated cells, where sperm spent  $94\pm2\%$  of their time in type 3 behaviour (Figure 3b;  $P=3*10^{-9}$  compared to control; Kruskal-Wallis with post-hoc comparison). Examination of the effect of stimuli on the characteristics of transitions between behaviours also showed that, as at pH 7.4,  $Ca^{2+}$ -mobilising treatments biased switching events towards types 2 and 3 behaviour (Figure 3c;  $P<0.005$ ; Chi-square). Dwell times for type 1 behaviour were not significantly altered by  $Ca^{2+}$ -mobilising stimuli at pH 8.5, but the effects of agonists on type 3 dwell times were more consistent and greater than at  $pH_o=7.4$ , with all four stimuli significantly extending the duration of type 3 behaviour compared to control cells (Figure 3f;  $P<0.001$ ; Kruskal-Wallis with post-hoc comparison). Most strikingly, in the presence of 4-AP, type 3 behaviour median dwell time was 56 s (compared to 3 s in

292 controls;  $P=10^{-16}$ , Kruskal-Wallis with post-hoc comparison; Figure 3f). Only thimerosal  
293 significantly increased type 2 dwell times at both  $pH_o=7.4$  and  $pH_o=8.5$  ( $P=1*10^{-10}$  and  
294  $P=1*10^{-5}$  respectively; Kruskal-Wallis with post-hoc comparison). Comparison of mean  
295 behaviour scores and % hyperactivated cells (CASA) for the 10 conditions assessed (control  
296 and four agonists, each at pH 7.4 and pH 8.5) showed a positive correlation between the two  
297 assessments of motility ( $R^2=0.5$ ;  $P=0.022$ ; Figure S8b).



## Discussion

By continuously observing ‘capacitated’, free-swimming human sperm, for a period of longer than 3 min, we have shown, for the first time, the occurrence of continuous switching between different types of motility, with the mean incidence being more than six transitions per minute. Even in control (unstimulated) cells, multiple behaviours, including hyperactivated-like motility, were seen in  $\approx 90\%$  of cells. For analysis of this switching, the intervening periods, where behaviour appeared consistent, were classified using four categories. Although tracks from behaviour types 1-3 resemble those referred to as activated, transitional and hyperactivated motility (Mortimer, 2000), these behaviours were derived by visual analysis and cannot be considered equivalent to categories defined by kinematic criteria. Type 4 behaviour where the flagellum arrested, typically in a ‘J’ shape, appears to be equivalent to the freeze and freeze-flex behaviours, which were observed in  $\approx 7\%$  of cells from normal semen (Burkman, 1984). In cells where analysis of FD was possible, it was clear that (i) visual analysis and FD identified the same behavioural transitions and (ii) for each recording, the mean FD was clearly and positively correlated with mean behaviour score ( $R^2=0.56$ ). We conclude that our analysis reliably identifies sperm behaviours and switching between them and that continuous behavioural switching is a normal activity of human sperm (at least for cells observed in vitro). Though some behaviour-time plots appeared to show oscillatory behaviour, autocorrelation analysis did not reveal significant periodicity.

The occurrence of repeated behavioural switching might be brought about by either, or both, of two different mechanisms. Changes in behaviour might be completely reliant on endogenous signalling activity, for instance,  $[Ca^{2+}]_i$  spikes and oscillations that have been described in human sperm (Harper et al, 2004; Aitken & McLaughlin, 2007; Sanchez-Cardenas et al, 2014; Mata-Martinez et al, 2018). These changes were observed in

immobilised cells and were correlated with changes in flagellar beating (Harper et al, 2004; Bedu-Addo et al, 2007). However, it is yet to be established whether these  $[Ca^{2+}]_i$  signals cause or are caused by the changes in flagellar beating. Alternatively, if mechanisms controlling sperm behaviour are sensitive to changes in the environment, such as pH, ionic environment, temperature and mechanical stimulation (including fluid flow), heterogeneity in the sperm's environment might present such 'stimuli' as it swims. Our data indicate that potential methodological artefacts related to movements of the stage and/or evaporation of the saline did not cause such effects (see Methods; Long Duration Sperm Tracking). However, the environment through which the sperm swims, even in the simple system used for our recordings, might be sufficiently spatially heterogenous that the sperm encounters a series of physico-chemical 'stimuli' due to its own movement. The complex environment encountered by the sperm in the female tract will almost certainly present such 'stimuli'. If behavioural switching is induced in this way, our observation that  $[Ca^{2+}]_i$  elevating stimuli prolong periods of type 3 behaviour (Figures 2f, 3f; see below) might indicate that  $Ca^{2+}$ -signalling resets the stimulus-sensitivity of transition into this type of motility. In either case, it appears likely that conditions/stimuli that result in behavioural switching in vitro will also occur in vivo.

Analysis of the effects of manoeuvres that elevate  $[Ca^{2+}]_i$  in human sperm indicate that motility kinematics (assessed by CASA) are directly related to the absolute level of  $[Ca^{2+}]_i$ , irrespective of the  $Ca^{2+}$  source mobilised, with the percentage of hyperactivated cells increasing as a function of  $[Ca^{2+}]_i$  (Achikanu et al, 2018). Exposure of cells to a range of stimuli which elevate  $[Ca^{2+}]_i$  significantly altered the relative abundance of the different behaviours. Consistent with a shift toward hyperactivated motility, the proportion of time spent in type 2 and type 3 behaviours markedly increased in cells exposed to  $Ca^{2+}$ -mobilising stimuli. Thimerosal and 4-AP were particularly potent in this regard. Thimerosal modifies

sperm behaviour by mobilising stored  $\text{Ca}^{2+}$  (Ho and Suarez, 2001; Alasmari et al., 2013b). Although 4-AP is often used as a  $\text{K}^+$  channel blocker, its effects on the  $\text{K}^+$ -permeable channels expressed in sperm are negligible (Tang et al, 2010; Mansell et al, 2014) and a significant aspect of its action on sperm  $[\text{Ca}^{2+}]_i$  and behaviour is likely to be its ability to mobilise stored  $\text{Ca}^{2+}$  (Gobet 1995; Grimaldi 2001; Baskhar 2008; Alasmari et al., 2013b Kasatkina, 2016), The potent effects of  $\text{Ca}^{2+}$ -store-mobilising stimuli on switching are consistent with their ability to persistently elevate  $[\text{Ca}^{2+}]_i$ , inducing prolonged hyperactivation in human sperm (Achikanu, et al., 2018; Alasmari, et al., 2013a,b). When cells were suspended in medium buffered at pH 8.5, some effects of  $\text{Ca}^{2+}$ -mobilising stimuli were greatly enhanced, particularly the prolongation of type 3 behaviour dwell-time by 4-AP. The effect of 4-AP on  $[\text{Ca}^{2+}]_i$  is strongly potentiated at pH=8.5 (Achikanu et al., 2018) and it is likely that this underlies the striking effect of the drug on behavioural switching under these conditions. Significantly, these potent hyperactivating stimuli had negligible effect on the proportion of cells in which type 3 (hyperactivated-like) behaviour was observed. Even under control conditions, type 3 behaviour (and behavioural switching) occurred in 90% of cells. Thus it appears that the well-characterised ability of these  $[\text{Ca}^{2+}]_i$ -mobilising compounds to increase the level of hyperactivation detected in population motility assays occurs not by recruitment of cells into a hyperactivated population, but by increasing the proportion of time that continuously-switching cells spend in type 3 (hyperactivated-like) motility, achieved primarily by extending the dwell-time of this behaviour.

If behavioural switching occurs in vivo, does it have adaptive value? Switching induced externally (by the sperm's sensitivity to heterogeneity in its environment; see discussion above) may simply be behavioural 'noise' that can be modulated by the cells signalling activity (such as elevation of  $[\text{Ca}^{2+}]_i$  as described here) but may not be functionally significant. However, observations of sperm interacting with the female tract and cumulus-

oocyte-complex suggest that switching of motility types may be of value. Pacey et al (1995) described an attach-detach cycle in the interaction of human sperm with epithelial cells isolated from the isthmic and ampullary sections of human oviducts, that may play an important role in migration to the fertilisation site. Detachment was associated with hyperactivated-type motility whereas cells that were attaching showed far more linear motility. Similarly, observation of sperm during the passage through the zona reveals alternation of low and high amplitude flagellar beats, which may facilitate penetration by alternation of ‘cutting’ and ‘thrusting’ forces (Bedford, 1998). Clearly further work is required to determine the nature and complexity of sperm behaviour in vivo and its significance.

**Authors' roles:** CA, JC and SMDS obtained samples and carried out the laboratory work. CA, SP and HG analysed the data. CA, JC, HG, LG, CLRB, SMDS and SJP contributed to writing of the manuscript.

**Funding:** This study was support by the Medical Research Council (MR/M012492/1). CA was in receipt of a scholarship from the Nigerian government (Tertiary Education Trust (TET) Fund).

**Conflict of interest:** C.L.R.B. was the editor-in-chief of Molecular Human Reproduction, has received lecturing fees from Merck and Ferring and is on the Scientific Advisory Panel for Ohana BioSciences. C.L.R.B. was also chair of the World Health Organization Expert Synthesis Group on Diagnosis of Male infertility (2012-2016). He is an editor of RBMO.

397 CLRB reports grants from the MRC, Gates Foundation and the Scottish Office during the  
398 conduct of the study; personal fees from Pharmasure, Ferring, Ohana, and RBMO outside the  
399 submitted work. CA, JC, HG, LG, SMDS and SJP declare no conflict of interest.

400

## References

- Achikanu C, Pendekanti V, Teague R, Publicover S. Effects of pH manipulation, CatSper stimulation and  $\text{Ca}^{2+}$ -store mobilization on  $[\text{Ca}^{2+}]_i$  and behaviour of human sperm. *Hum Reprod* 2018;33: 1802-1811.
- Aitken RJ, McLaughlin EA. Molecular mechanisms of sperm capacitation: progesterone-induced secondary calcium oscillations reflect the attainment of a capacitated state. *Society of Reproduction and Fertility supplement* 2007;63: 273-293.
- Alasmari W, Barratt C, Publicover S, Whalley K, Foster E, Kay V, Martins da Silva S, Oxenham S. The incidence and clinical significance of defects in calcium signalling pathways mediating human sperm hyperactivation in donors and sub fertile patients. *Human Reproduction* 2013a;28: 866-876.
- Alasmari W, Costello S, Correia J, Oxenham SK, Morris J, Fernandes L, Ramalho-Santos J, Kirkman-Brown J, Michelangeli F, Publicover S, Barratt C.  $\text{Ca}^{2+}$  signals generated by CatSper and  $\text{Ca}^{2+}$  regulate different behaviours in human sperm. *J Biol Chem* 2013b;288: 6248-6258.
- Bhaskar A, Subbanna PK, Arasan S, Rajapathy J, Rao JP, Subramani S. 4-aminopyridine-induced contracture in frog ventricle is due to calcium released from intracellular stores. *Indian J Physiol Pharmacol* 2008;52: 366-374.
- Alvarez L, Dai L, Friedrich BM, Kashikar ND, Gregor I, Pascal R, Kaupp UB. The rate of change in  $\text{Ca}^{2+}$  concentration controls sperm chemotaxis. *J Cell Biol* 2012; 5: 653-663
- Bedford JM. Mammalian fertilization misread? Sperm penetration of the eutherian zona pellucida is unlikely to be a lytic event. *Biol Reprod* 1998;59: 1275-1287.

424 Bedu-Addo K, Barratt CL, Kirkman-Brown JC, Publicover SJ. Patterns of  $[Ca^{2+}]_i$   
 425 mobilization and cell response in human spermatozoa exposed to progesterone. *Dev Biol*  
 426 2007;302: 324-332.  
 427 Brown SG, Costello S, Kelly MC, Ramalingam M, Drew E, Publicover SJ, Barratt CLR, Da  
 428 Silva SM. Complex CatSper-dependent and independent  $[Ca^{2+}]_i$  signalling in human  
 429 spermatozoa induced by follicular fluid. *Hum Reprod* 2017;32: 1995-2006.  
 430 Burkman LJ. Characterisation of hyperactivated motility by human spermatozoa during  
 431 capacitation: comparison of fertile and oligozoospermic sperm populations. *Arch Androl*  
 432 1984;13: 153.  
 433 Burkman LJ. Characterization of hyperactivated motility by human spermatozoa during  
 434 capacitation: comparison of fertile and oligozoospermic sperm populations. *Arch Androl*  
 435 1984;13: 153-165.  
 436 Carlson AE, Westenbroek RE, Quill T, Ren D, Clapham DE, Hille B, Garbers DL, Babcock  
 437 DF. CatSper1 required for evoked  $Ca^{2+}$  entry and control of flagellar function in sperm. *Proc*  
 438 *Natl Acad Sci U S A* 2003;100: 14864-14868.  
 439 Cooper G, Overstreet J, Katz D. The motility of rabbit spermatozoa recovered from the  
 440 female genital tract. *Gam Res* 1979;2: 35-42.  
 441 Gabel CP, Carroll J, Harrison K. Sperm motility is enhanced by Low Level Laser and Light  
 442 Emitting Diode photobiomodulation with a dose-dependent response and differential effects  
 443 in fresh and frozen samples. *Laser Ther* 2018;27: 131-136.  
 444 Gaetano J. Holm-Bonferroni Sequential Correction: An EXCEL calculator (1.2).  
 445 <https://wwwresearchgatenet/publication/242331583> 2013.  
 446 Gobet I, Lippai M, Tomkowiak M, Durocher Y, Leclerc C, Moreau M, Guerrier P. 4-  
 447 aminopyridine acts as a weak base and a  $Ca^{2+}$  mobilizing agent in triggering oocyte meiosis

448 reinitiation and activation in the Japanese clam *Ruditapes philippinarum*. *Int J Dev Biol*  
 449 1995;39: 485-491.

450 Grimaldi M, Atzori M, Ray P, Alkon DL. Mobilization of calcium from intracellular stores,  
 451 potentiation of neurotransmitter-induced calcium transients, and capacitative calcium entry  
 452 by 4-aminopyridine. *J Neurosci* 2001;21: 3135-3143.

453 Harper CV, Barratt CL, Publicover SJ. Stimulation of human spermatozoa with progesterone  
 454 gradients to simulate approach to the oocyte. Induction of  $[Ca^{2+}]_i$  oscillations and cyclical  
 455 transitions in flagellar beating. *J Biol Chem* 2004;279: 46315-46325.

456 Ho HC, Suarez SS. An inositol 1,4,5-trisphosphate receptor-gated intracellular  $Ca^{2+}$  store is  
 457 involved in regulating sperm hyperactivated motility. *Biol Reprod* 2001;65: 1606-1615.

458 Ho K, Wolff CA, Suarez SS. CatSper-null mutant spermatozoa are unable to ascend beyond  
 459 the oviductal reservoir. *Reprod Fertil Dev* 2009;21: 345-350.

460 Johnson LL, Katz, D.F., Overstreet. J.W. The movement characteristics of rabbit  
 461 spermatozoa before and after activation. *Gamete Res* 1981;4: 275-282.

462 Kasatkina LA. 4-Aminopyridine sequesters intracellular  $Ca^{2+}$  which triggers exocytosis in  
 463 excitable and non-excitable cells. *Scientific reports* 2016;6: 34749.

464 Katz DF, Yanagimachi R. Movement characteristics of hamster spermatozoa within the  
 465 oviduct. *Biol Reprod* 1980;22: 759-764.

466 Mansell SA, Publicover SJ, Barratt CL, Wilson SM. Patch clamp studies of human sperm  
 467 under physiological ionic conditions reveal three functionally and pharmacologically distinct  
 468 cation channels. *Mol Hum Reprod* 2014;PMID: 24442342: 392-408.

469 Mata-Martinez E, Darszon A, Trevino CL. pH-dependent  $Ca^{2+}$  oscillations prevent untimely  
 470 acrosome reaction in human sperm. *Biochem Biophys Res Commun* 2018;497: 146-152.

471 Mortimer ST. CASA - practical aspects. *J Androl* 2000;21: 515-524.



472 Mortimer ST, Swan MA. Kinematics of capacitating human spermatozoa analysed at 60 Hz.  
 473 *Hum Reprod* 1995;10: 873-879.

474 Mortimer ST, Swan MA, Mortimer D. Fractal analysis of capacitating human spermatozoa.  
 475 *Hum Reprod* 1996;11: 1049-1054.

476 Pacey AA, Davies N, Warren MA, Barratt CL, Cooke ID. Hyperactivation may assist human  
 477 spermatozoa to detach from intimate association with the endosalpinx. *Hum Reprod* 1995;10:  
 478 2603-2609.

479 Pacey AA, Ladbroke MB, Barratt CL, Cooke ID. The potential shortcomings of measuring  
 480 hyperactivated motility by computer-aided sperm analysis when sperm motion is multiphasic.  
 481 *Hum Reprod Update* 1997;3: 185-193.

482 Publicover S, Harper CV, Barratt C.  $[Ca^{2+}]_i$  signalling in sperm--making the most of what  
 483 you've got. *Nat Cell Biol* 2007;9: 235-242.

484 Sanchez-Cardenas C, Montoya F, Navarrete, FA, Hernandez-Cruz A, Corkidi G, Visconti PE,  
 485 Darszon A (2018) Intracellular  $Ca^{2+}$  threshold reversibly switches flagellar beat off and on.  
 486 *Biol Reprod* doi: 10.1093/biolre/iox132.

487 Sanchez-Cardenas C, Servin-Vences MR, Jose O, Trevino CL, Hernandez-Cruz A, Darszon  
 488 A. Acrosome reaction and  $Ca^{2+}$  imaging in single human spermatozoa: new regulatory roles  
 489 of  $[Ca^{2+}]_i$ . *Biol Reprod* 2014;91: 67.

490 Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.  
 491 Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J. Y. Tinevez, D. J. White, V. Hartenstein, K.  
 492 Eliceiri, P. Tomancak and A. Cardona (2012). Fiji: an open-source platform for biological-  
 493 image analysis. *Nat Methods* 2012; 9: 676-682.

494 Shahar S, Wiser A, Ickowicz D, Lubart R, Shulman A, Breitbart H. Light-mediated activation  
 495 reveals a key role for protein kinase A and sarcoma protein kinase in the development of  
 496 sperm hyper-activated motility. *Hum Reprod* 2011;26: 2274-2282.

Suarez SS. Control of hyperactivation in sperm. *Hum Reprod Update* 2008;14: 647-657.

Tang QY, Zhang Z, Xia XM, Lingle CJ. Block of mouse Slo1 and Slo3 K<sup>+</sup> channels by

CTX, IbTX, TEA, 4-AP and quinidine. *Channels (Austin)* 2010;4: 22-41.

# **Figure 1. Behavioural switching in a free-swimming human sperm under control**

**conditions (pH<sub>o</sub>=7.4).** (a) Variation in behaviour type (categorised visually as type1, type 2,

type 3 or type 4; see Figure S1) of a single sperm over a period of 190 s. (b) Variation in

fractal dimension (FD) over time (black trace) overlaid with the visually categorised


behaviour types (red trace). Visual analysis and FD show good agreement, with no visually

identified behavioural transitions that are not confirmed by FD. (c) Track of the same cell,


colour coded to display variation in the fractal dimension (FD); ( $1 < FD \leq 1.2$  (dark blue);

$1.2 < FD \leq 1.4$  (light blue);  $1.4 < FD \leq 1.6$  (green);  $1.6 < FD \leq 1.8$  (yellow);  $1.8 < FD \leq 2.0$  (red). Axes


show distance in mm.



**Figure 2. Characteristics of behavioural switching.** (a) Dwell times (period during which a single behaviour type occurred) for types 1 (light blue, n=164 events), 2 (green, n=79), 3 (red, n=107) and 4 (black, n=15) behaviours in cells control cells (pH 7.4). Plots show median and interquartile range (box) and maximum/minimum values (whiskers). (b) Mean % time spent in each behaviour; type 1 (light blue), type 2 (green), type 3 (red) and type 4 (black) under control conditions (n=18 cells) and in the presence of 300 nM P4 (n=18 cells), 1% FF (n=17 cells), 2 mM 4-AP (n=21 cells) and 1 mM thimerosal (n=17 cells). (c) Relative frequencies of transitions into type 1 (light blue), type 2 (green), type 3 (red) and type 4 (black) behaviours under control conditions (n=347 transitions) and in the presence of progesterone (P4; 300 nM; n=420), human follicular fluid (FF nM, 1%; n=353), 4-aminopyridine (4-AP, 2 mM, n=211) and thimerosal (1 mM, n=115). Asterisks indicate significant difference from control; \*\*\*=P<0.005. (d-f) Dwell times for type 1 (panel 'd'; blue), type 2 (panel 'e'; green) and type 3 (panel 'f'; red) behaviours under control conditions and in the presence of 300 nM progesterone (P4), 1% human follicular fluid (FF), 2 mM 4-aminopyridine (4-AP) and 1 mM thimerosal. Plots show median and interquartile range (box) and maximum/minimum values (whiskers) of 51-167 events. Asterisks indicate significant difference from control; \*\*=P<0.01; \*\*\*=P<0.001; \*\*\*\*=P<0.0001.



**Figure 3. Characteristics of behavioural switching at  $\text{pH}_0=8.5$ .** (a) Mean switching rate ( $\pm$ s.e.m.) for cells incubated at  $\text{pH}_0=7.4$  (grey bars) and  $\text{pH}_0=8.5$  (red bars) under control conditions ( $n=18$  cells) and in the presence of 300 nM P4 ( $n=18$  cells), 1% FF ( $n=17$  cells), 2 mM 4-AP ( $n=21$  cells) and 1 mM thimerosal ( $n=17$  cells). (b) Mean % time spent in each behaviour at  $\text{pH}_0=8.5$ ; type 1 (light blue), type 2 (green), type 3 (red) and type 4 (black) under control conditions ( $n=18$  cells) and in the presence of 300 nM P4 ( $n=15$  cells), 1% FF ( $n=18$  cells), 2 mM 4-AP ( $n=20$  cells) and 1 mM thimerosal ( $n=16$  cells). (c) Relative frequencies of transitions into type 1 (light blue), type 2 (green), type 3 (red) and type 4 (black) behaviours at  $\text{pH}_0=8.5$  under control conditions ( $n=220$  transitions) and in the presence of progesterone (P4; 300 nM;  $n=194$ ), human follicular fluid (FF nM, 1%;  $n=151$ ), 4-aminopyridine (4-AP, 2 mM,  $n=47$ ) and thimerosal (1 mM,  $n=63$ ). (d-f) Dwell times at  $\text{pH}_0=8.5$  for type 1 (panel 'd'; blue), type 2 (panel 'e'; green) and type 3 (panel 'f'; red) behaviours under control conditions and in the presence of 300 nM progesterone (P4), 1% human follicular fluid (FF), 2 mM 4-aminopyridine (4-AP) and 1 mM thimerosal. Plots show median and interquartile range (box) and maximum/minimum values (whiskers) of 10-110 events (except thimerosal type 1 where  $n=2$ ). Asterisks indicate significant difference from control;  $*=P<0.05$ ;  $**=P<0.01$ ;  $***=P<0.001$ ;  $****=P<0.0001$ .



569 **Supplementary video legends**

570 **Video 1. Example of repeated behavioural switching.** In this cell type, 1 behaviour occurs  
571 at the start of the video and also for the periods 3-5.5 s, 11.5-14 s and 17.5-20 s as shown by  
572 the time stamp (seconds) at the top left of the image. Periods of type 3 behaviour occur at  
573 approximately 2-3 s, 10-11.5 s and 14-17.5 s. Between 5.5 and 10 s both type 3 and (briefly)  
574 type 2 behaviours occur. Total duration= 20 s. Frame dimensions = 230 mm \* 225 mm

575 **Video 2. Example of a cell showing type 2 behaviour.** Total duration= 6 s. Frame  
576 dimensions = 230 mm \* 240 mm

577 **Video 3. Example of a cell showing type 4 (arrested) motility.** This cell shows periods of  
578 type 4 motility (where the flagellum arrests in a J shape) interspersed by brief periods (< 1 s)  
579 of flagellar beating. Total duration= 5.9 s. Frame dimensions = 205 mm \* 180 mm.

580

**Table 1. Characteristics of the motility types identified by visual analysis of videos.** Values shown for typical ALH (amplitude of lateral head displacement) show range of values obtained from examination of 15-20 tracks of each type and are descriptive, not definitive. n/a indicates not applicable.

motility type	characteristics	progressive/non-progressive	typical ALH
1	low amplitude flagellar beat, symmetric or occasionally weakly asymmetric causing curved or circular path	progressive	≈2-4 μm
2	intermediate amplitude beat, symmetric or slightly asymmetric (causing circling)	progressive	≈3-8 μm
3	high-amplitude highly asymmetric beat	non-progressive, continuous turning or tumbling	≥6 μm
4	tight bending of the midpiece producing a 'J'-shape or coil	non-progressive arrested	n/a

**Table 2. Incidence of behavioural switching.** Data show proportion of cells in which behavioural switching was observed (number of cells showing switching/total number of cells analysed) at pH<sub>o</sub>=7.4 and pH<sub>o</sub>=8.5 under control conditions and in the presence of 300 nM progesterone (P4), 1% human follicular fluid (FF), 2 mM 4-aminopyridine (4-AP) and 1 μM thimerosal. The bottom row of the table (P) shows results of chi-square test to compare between pH values (corrected for multiple comparisons) under each condition.

pH <sub>o</sub>	treatment				
	control	P4	FF	4-AP	thimerosal
7.4	16/18	16/18	16/17	20/21	14/19
8.5	16/18	13/15	14/18	10/20	11/16
P	1	1	0.67	0.01	1